

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

#4 Attachment
09/6224/9

20



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 236 429 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: 07.04.93 (51) Int. Cl.⁵: C12N 15/52

(21) Application number: 86905574.9

(22) Date of filing: 26.08.86

(86) International application number:
PCT/US86/01759

(87) International publication number:
WO 87/01391 (12.03.87 87/06)

(54) SYSTEM FOR BIOTIN SYNTHESIS.

(30) Priority: 26.08.85 US 769849
12.08.86 US 893042

(43) Date of publication of application:
16.09.87 Bulletin 87/38

(45) Publication of the grant of the patent:
07.04.93 Bulletin 93/14

(64) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(56) References cited:

CHEMICAL ABSTRACTS, vol. 93, no. 7, 18th August 1980, page 480, abstract no. 65744k, Columbus, Ohio, US; A. MUKHERJEE et al.: "Construction and characterization of a recombinant plasmid containing the entire biotin genes of E. coli K12", & PLASMIDS TRANSPOSONS, [PROC. ANNU. SYMP. SCI. BASIS MED.], 4TH 1979 (PUB. 1980), 379-86

(73) Proprietor: Amgen
1900 Oak Terrace Lane
Thousand Oaks California 91320(US)

(72) Inventor: FISHER, Eric, F.
711 Flaming Star Drive
Thousand Oaks, CA 91360(US)

(74) Representative: Brown, John David et al
FORRESTER & BOEHMERT Franz-
Joseph-Strasse 38
W-8000 München 40 (DE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

MOLEC. GEN. GENET., v 1. 166, 1978, pages 305-312; G. COHEN et al.: "Isolation and characterization of a ColE1 plasmid containing the entire biotin gene cluster of Escherichia coli K12"

J. MOL. BIOL., vol. 148, 1981, pages 63-76, Academic Press Inc., Ltd, London, GB; A. SANCAR et al.: "Identification of the uvrB gene product"

GENE, 35, August 1985, Howard et al, "Nucleotide Sequence of the biA Gene Encoding the Biotin Operon Holoenzyme Synthetase Functions of Escherichia Coli", pp. 321-331

JOURNAL OF BACTERIOLOGY, 112(3), December 1972, PAI, "Mutant of Escherichia Coli with Depressed Levels of the Biotin Biosynthetic Enzymes" pp. 1280-1287

JOURNAL OF BACTERIOLOGY, 112(2), November 1972, Cleary et al, "Deletion and Complementation Analysis of the Biotin Gene Cluster of Escherichia Coli, pp. 830-839.

JOURNAL OF BACTERIOLOGY, 14 (2), August 1980, Barker et al, "Use of bio-lac Fusion Strains to Study Regulation of Biotin-Biosynthesis in Escherichia Coli", pp. 789-800

US 4 563 426, (YAMADA) 07 January 1986, Column 1, l. 58-68

Ja 0.060.996, (NIPPON ZEON KK), 11 April 1983, Abstract

D scription

This is a continuation-in-part of co-pending U.S. Patent Application Serial No. 769,849, filed August 26, 1985.

Background

The present invention relates in general to systems for the microbial production of biotin and in particular to systems wherein at least a part of the biotin operon is present on a plasmid within a biotin retention-deficient mutant host cell.

Biotin, also known as vitamin H, is probably an essential component of all cells. Some microorganisms, including baker's yeast, and all animals (except the protozoan *Tetrahymena*) are unable to synthesize biotin effectively and must therefore obtain biotin from their environment in order to survive.

Despite its usefulness in promoting the growth of baker's yeast and as a human and animal food additive, biotin is very expensive to manufacture by presently available, chemical synthetic methods. Furthermore, although beet molasses (containing 0.015-0.15 μ grams of biotin per gram) or other natural sources of biotin may be used to supplement synthetic biotin, there exists a need for other sources.

Due to the ready availability of information regarding the genetic constitution of certain microorganisms which have been reported to contain relatively high concentrations of biotin, a capability for performing genetic manipulations on those microorganisms has developed. It has been reported, for example, that certain chromosomal genes which encode enzymes of the pathway for biotin synthesis may be isolated, amplified and reinserted into host cells of the bacterium *Escherichia coli* (*E. coli*).

More specifically, Mukherjee, et al. in "Plasmids and Transposons" Stutard, et. al., eds., Academic Press, New York (1980), 379-386 reported isolation of the biotin operon of the *E. coli* K-12 strain from a transducing bacteriophage by means of *EcoRI* enzyme digestion. A restriction fragment was inserted into a DNA plasmid (pMB8) which was used to transform *E. coli* host cells to provide multiple "extra" copies of the biotin operon genes in these hosts. Mukherjee, et. al., however fails to teach or even suggest the use of a biotin retention-deficient mutant genotype host cell. Although enhancement of excretion over a biotin prototroph ("wild type") was reported, this recombinant system has not been applied to large scale commercial production of biotin by fermentation of transformed host cells.

SUMMARY OF THE INVENTION

According to the present invention there is provided a system for the production of biotin comprising: a microorganism having a *birA*⁻ genotype and plasmid DNA, within said cell, encoding at least one gene product of the biotin operon or a functional homolog thereof.

Another aspect of the present invention comprises a system for the production of biotin comprising: a microorganism having a (*bio*⁻, *birA*⁻, *bioR*⁻) genotype; and plasmid DNA, within said cell, wherein said plasmid DNA encodes the biotin operon and wherein said plasmid DNA is *uvrB*⁻.

The present invention further relates to a system for the production of biotin comprising: a microorganism having a (*bio*⁻, *birA*^{ts}, *bioR*⁻) genotype; and plasmid DNA encodes the biotin operon and wherein said plasmid DNA is *uvrB*⁻.

There is further provided by the present invention a method for the enhanced production of biotin which comprises transforming a microorganism having a *birA*⁻ genotype with a plasmid DNA encoding a gene product of the biotin operon or a functional homolog thereof.

According to the present invention there is still further provided a method of converting desthiobiotin to biotin comprising the steps of: culturing a host microorganism, having a biotin retention-deficient mutant genotype and having plasmid DNA encoding a *bioB* gene product of the biotin operon or a functional homolog thereof, in a medium containing desthiobiotin.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a flowchart of the biotin biosynthetic pathway;

Fig. 2 is a schematic illustration of the *bioR* and *birA* gene functions;

Fig. 3 is a partial restriction map of the *bio* (A, B, F, C, D) operon and the adjacent *uvrB* locus on the *E. coli* chromosome;

Fig. 4 is a schematic illustration of the construction of the intermediate plasmid 322PstI according to the present invention;

Fig. 5 is a schematic illustration of the construction of a bioD gene restriction fragment according to the present invention;

Fig. 6 is a schematic illustration of the construction of the intermediate plasmid pBAL4 according to the present invention;

Fig. 7 is a schematic illustration of the construction of the biotin operon-containing plasmid pBP5 according to the present invention;

Fig. 8 is a schematic illustration of the construction of the plasmids pKA5 and pKH4 according to the present invention; and

Fig. 9 is a schematic illustration of the construction of the plasmid pBF1 according to the present invention.

DETAILED DESCRIPTION

As used herein, the term "biotin retention-deficient mutant genotype" refers to a lesion in birA gene, that produces a change in the birA gene resulting in a decrease in the activity of the birA gene product, that is, a mutation at the birA locus giving rise to a diminished capacity for adenylation of biotin and is hereinafter referred to as birA⁻. A class of preferred lesions in birA comprise lesions which will render the activity of the enzyme dependant on temperature, that is a temperature sensitive birA gene (birA^{TS}). Such birA^{TS} mutants decrease the birA function as the temperature of the system increases.

Preferred host cells include biotin-requiring strains (genotype: bio⁻) strains deficient in the repressor of the biotin operon (genotype: bioR⁻) and biotin-requiring strains deficient in the repressor function (genotype: bio⁻, bioR⁻). The most preferred host cells are bioR⁻ strains.

The term "functional homolog of a gene product of the biotin operon" refers to a polypeptide which has the same function as, but which may have the same amino acid sequence as or different in amino acid sequence from, the gene product. Such functional homologs include, for example, polypeptide products of allelic variations of the genes of the biotin operon; analogs and fragments of these polypeptides; and synthetic polypeptides which may be dissimilar in primary structure (amino acid sequence) but which share secondary structures that allow them to possess biological and immunological activities of gene products of the biotin operon [Kaiser, et al., Science, 223, 249-255 (1984)].

A flow chart of biotin biosynthesis is presented in Fig. 1. Six enzymes involved in biotin biosynthesis have been assigned to six genetic loci: bioA, bioB, bioC, bioD, bioF and bioH. Specific reactions catalyzed by bioA, bioB, bioD and bioF gene products have been characterized. Cofactors and substrates for each of these reactions, with the exception of a sulfur atom donor in the last enzymatic step, have been identified. Although the functions of the bioC and bioH gene products have not been characterized due to limitations of crossfeeding studies (wherein biotin-deficient strains obtain survival only by utilizing biotin or by utilizing biosynthetic precursors of biotin excreted by cells with which they are co-cultured), these loci have been identified as essential by genetic complementation.

The six genes coding for biotin synthetic enzymes are located in two regions of the *E. coli* chromosome. Five of the six genes (bioA, bioB, bioC, bioD and bioF) are contained in a bidirectionally transcribed operon mapped at 17 minutes. BioH is located at 74 minutes. The locations of the genes of the biotin operon and of two other genetic functions which impinge on the biotin biosynthetic pathway, bioR and birA, are given in Table I.

TABLE I

Genetic Locus	Location on <i>E. coli</i> Chromosome	Proposed Function
<u>bioA</u>	17 min	Synthetic enzyme
<u>bioB</u>	17 min	Synthetic enzyme
<u>bioC</u>	17 min	Synthetic enzyme
<u>bioD</u>	17 min	Synthetic enzyme
<u>bioF</u>	17 min	Synthetic enzyme
<u>bioH</u>	74 min	Synthetic enzyme
<u>bioR</u>	88 min	Transcription repressor
<u>birA</u>	88 min	Utilization of biotin

Control of biotin synthesis in *E. coli* is effected at the transcriptional level. After biotin is synthesized, it is adenylylated by a product of a gene at a locus designated birA, to form biotinyl-5'-adenylate as illustrated

in Fig. 2. A biotin repressor protein, identified as a product of the *bioR* locus may also bind to biotinyl-5'-adenylate to increase 25-fold the affinity of the *bioR* gene product for a *bio* operator. Howard, et. al., *Gene*, 35, 321-331, (1985), have disclosed that the *birA* function and the *bioR* function are effected by the same protein.

The *bio* operator is situated between the *bioA* structural gene and the *bioB* structural gene, as depicted in Fig. 3. The *bio* operator overlaps both the *bioA* gene promoter and the *bioB* gene promoter. The *bioR* gene product may terminate transcription by binding to the *bio* operator and excluding RNA polymerase from these two divergent promoters.

Biotinyl-5'-adenylate is also a substrate for what is believed to be a third function of the *birA* gene product, biotin holoenzyme synthetase. Biotin holoenzyme synthetase transfers biotin to acetyl-CoA carboxylase. Acetyl-CoA carboxylase catalyzes a critical step in fatty acid synthesis, which is essential for viability. This implies that a complete elimination of *birA* activity at the initiation of the fermentation would be lethal. Therefore, at the initiation of the fermentation, it is necessary that sufficient *birA* activity be present to support growth of the cells. Such *birA* activity is readily ascertained by one of ordinary skill in the art. Upon completion of the fermentation it is preferred that *birA* activity is substantially diminished and most preferably eliminated. By employing a *birA^{TS}* gene, it is possible to regulate the *birA* function by controlling the temperature of the fermentation system. Therefore as the temperature of the system is increased, the *birA* function of the cell is decreased.

A genetic locus which maps adjacent to the *bioD* locus is denominated *uvrB*. The *uvrB* gene has no function in biotin physiology, but acts in some way to protect *E. coli* cells from ultraviolet radiation as reported by Sancar, et al., *Cell*, 28, 523-530 (1982). Three RNA molecules are transcribed from the *uvrB* locus, one of which may interact with RNA polymerase A. Therefore, if the *uvrB* gene were multiplied, this interaction may be lethal for an *E. coli* cell. For this reason, prior to increasing the copy number of a plasmid containing a piece of DNA from the region of the *E. coli* chromosome which includes the biotin operon, *uvrB* functions should preferably be eliminated.

In addition to preferably employing a *birA^{TS}* gene, it is preferred to employ a plasmid having a high copy number, and most preferably a plasmid that exhibits a moderate increase in copy number (40 to 200) upon temperature induction. Such plasmids are described in European Patent Application No. 136,490 and are hereinafter referred to as temperature sensitive plasmids. Therefore when utilizing such temperature sensitive plasmids, it is possible upon increasing the temperature of the reaction to moderately increase the copy number and gene dosage while maintaining cell vitality and decreasing the *birA* function, thereby resulting in a system capable of producing surprisingly high yields of biotin. The following Examples serve to further illustrate embodiments of the present invention. Although the *birA⁻* strains of Barker, et al., *J. Bacteriol.*, 143, 789-800 (1980) and Campbell, et al., *Proc. Natl. Acad. Sci. (USA)*, 69, 676-680 (1972) are employed in the Examples, other biotin retention deficient strains such as for example the *E. coli* strain P48 reported in Pai, *J. Bacteriol.*, 112, 1280-1287 (1972), may also be employed.

Example 1

As illustrated in Fig. 4, a first plasmid, designated pLC2523 (deposited August 23, 1985 as Deposit No. A.T.C.C. 53237 with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland) and known to contain the biotin operon (see e.g., Sancar et al., *J. Mol. Biol.*, 148, 63-76, (1981)), and a second plasmid, designated pBR322 (ATCC No. 37017) were digested with *Pst*I and joined by T4 DNA ligase. The mixture was then transformed, according to the procedure of Hanahan, *J. Mol. Biol.*, 166, 557-580 (1983), into bacterial cells of the biotin auxotrophic strain SA291 (*bio⁻*, *bioR⁺*, *birA⁺*) (deposited August 23, 1985 with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland, as Deposit No. A.T.C.C. 53236). Colonies were selected on L agar plates containing tetracycline (12 mg/L), which killed any cells that did not contain plasmids having the tetracycline-resistance segment of pBR322. The selected colonies were screened for ampicillin sensitivity, an indication that a *Pst*I digestion fragment from pLC2523 had been inserted into the *Pst*I site of the ampicillin-resistance segment of pBR322, thereby rendering it incapable of conferring resistance.

Restriction fragments of plasmids conferring tetracycline resistance were separated by gel electrophoresis and examined for the presence of fragments having the expected lengths of the biotin operon. In this way it was determined that a plasmid designated 322PST1 contained the biotin operon linked to a tetracycline-resistance marker. However, in this plasmid the *bioD* gene was found not to be intact for it would not complement biotin auxotrophic *E. coli* strain SA291 when grown in the absence of biotin.

As illustrated in Fig. 5, in order to obtain the remainder of the *bioD* gene, pLC2523 was cleaved with *Nco*I to produce a larger fragment and a smaller fragment (4.4 kilobases in length) which were separated by

gel electrophoresis. The smaller fragment was extracted from the gel and both ends of the smaller fragment were digested with the exonuclease BAL31.

In the BAL31 digestion, 30 µg of the restriction fragment was dissolved in 450 µL of BAL31 nuclease buffer containing 0.25 mg/mL of bovine serum albumin. A 200 µL portion was treated with 2 units/µL of BAL31 at 30°C. Samples were withdrawn and phenol extracted at 2.5 min, 5.5 min and 10 min. After the extraction and ethanol precipitation, an aliquot of each time point sample was analyzed by electrophoresis through a 0.5% (w/v) agarose gel. The three time point samples were pooled. The shortened fragments thus obtained were further cleaved with BglII and PstI. Due to the presence of two BglII sites and one PstI site in the NcoI fragment of pLC2523, four types of fragments were expected. Of these four types of fragments, one was expected to contain the remainder of the bioD gene and to have both a blunt, BAL31-digested end and a cohesive PstI-digested end.

The plasmid pBR329, [the complete nucleotide sequence of which is published in Covarrubias, et al., Gene, 17, 79 (1982), which is incorporated by reference herein] was digested with both PvuII (which cleaves pBR329 in a chloramphenicol-resistance segment, to produce a blunt end) and with PstI (which cleaves pBR329 at a site in an ampicillin-resistance segment) to obtain two pieces which were separated by gel electrophoresis. The larger of the pieces (containing a tetracycline resistance segment and an origin of replication) was mixed with the four types of fragments produced by the BglII and PstI digestion of the 4.4 kb NcoI fragment described above in the presence of T4 DNA ligase. As shown in Fig. 6, only those fragments containing the remainder of the bioD gene had the combination of blunt and PstI-digested ends required to join with the larger PvuII/PstI fragment from pBR329 to form a cyclic plasmid designated pBAL4.

Bacteria of strain SA291 were transformed with the products of the ligation with the larger fragment of pBR329. Colonies were selected for tetracycline resistance, screened for ampicillin sensitivity and screened for chloramphenicol sensitivity. The lengths of the inserts in various plasmids were determined by restriction endonuclease analysis.

As shown in Fig. 7, the plasmids 322PstI and pBAL4 were separately digested with PstI. These digests were combined in a ligation reaction, using T4 DNA ligase. The resulting mixture was used to transform the cells of strain SA291. Colonies were selected for a combination of growth in the absence of biotin and growth in the presence of 12 mg/mL tetracycline. Presence of the complete bio operon was confirmed by retransformation of plasmid-free SA291 in conjunction with restriction endonuclease digestion analysis. A resulting plasmid, designated pBP5, contained all of the genes of the biotin operon: genes bioA, bioB, bioF, bioC and the portion of the bioD gene upstream of the PstI site derived from 322PstI and the portion of the bioD gene downstream of the PstI site derived from pBAL4.

Next, the temperature sensitive copy number plasmid pCFM 526 was digested with EcoRI and resealed with ligase to produce pCFM 526ΔE4, which lacked the P_L promoter contained in pCFM 526. Plasmid pCFM526 had been constructed as described in Morris, published European Patent Application No. 136,490, from plasmid pCFM414 (ATCC No. 40,076).

As shown in Fig. 8, plasmid pCFM526ΔE4 and the plasmid pBP5 were separately digested with HindIII. The fragments were ligated and used to transform SA291. Colonies were selected for ampicillin resistance and the ability to grow in the absence of biotin. A plasmid designated pKA5 was isolated. This plasmid contained the five genes of the bio operon linked to a temperature inducible origin of replication.

Example 2

As further shown in Fig. 8, another plasmid was also constructed in a fashion analogous to the construction of pKA5 described in Example 1, but with the substitution of a plasmid designated pCFM1036NS, which contains a kanamycin resistance segment, for pCFM526ΔE4. Colonies were therefore selected for kanamycin resistance instead of ampicillin resistance to obtain cells bearing a plasmid pKH4.

EXAMPLE 3

A HindIII fragment from pBP5 was treated with BAL31 and the mixture was ligated into HpaI-cut pCFM526ΔE4. Colonies were selected for biotin production, ampicillin resistance, and tetracycline sensitivity. Three plasmids, pBA2, pBA4 and pBA6 were obtained from this selection.

EXAMPLE 4

As illustrated in Fig. 9, the plasmids pBP5 and pCFM526 were cut with NcoI and HindIII. The ligation product of these digests was transformed into an E. coli strain AM7 containing plasmid pMW1 (A.T.C.C. No.

39933) harboring a gene for the temperature sensitive repressor C^{857} . In this construction, designated pBF1, the *bioB* gene is placed under the control of the P_L promoter. Therefore, this construction is useful for converting desthiobiotin into biotin by means of the *bioB* gene product, biotin synthetase.

5 EXAMPLE 5

The plasmid pLC2523 was digested with HindIII and NcoI. The plasmid pCFM526 was similarly cut. A ligation product of these two digests, designated pAHN203, was transformed into cells of a bacterial strain containing the temperature-sensitive repressor of bacteriophage λ (C^{857}). The plasmid pCFM526 contains the P_L promoter of bacteriophage λ . A gene or genes inserted downstream from this locus are controlled by this promoter. The promoter activity is regulated by repressor C^{857} . Therefore, when the temperature is raised, the repressor function is eliminated, the promoter is activated and the desired gene products are expressed. See e.g., Morris, supra. In pAHN203 the *bioA* gene is under P_L control. The plasmid pAHN203 is combined with pBF1 to produce a plasmid which produces biotin under P_L control.

The following buffers were employed in the Examples. A high salt buffer comprising: 75mM NaCl; 50mM Tris-HCl, pH 7.6; 10mM MgCl₂; and 5mM dithiothreitol. A medium salt buffer comprising: 37.5mM NaCl; 30mM Tris-HCl, pH 7.6; 10mM MgCl₂; and 5mM dithiothreitol. A low salt buffer comprising: 10mM Tris-HCl, pH 7.6; 10mM MgCl₂; 20mM KCl; and 5mM dithiothreitol. A ligase buffer comprising: 50mM Hepes, pH 7.5; 10mM MgCl₂; 5mM dithiothreitol; and 0.4mM adenosine triphosphate. A polynucleotide kinase buffer comprising: 50mM Tris-HCl, pH 7.6; 10mM MgCl₂; 1mM spermidine; 5mM dithiothreitol; and 0.1 mM ethylenediamine tetraacetic acid (EDTA). A BAL31 nuclease buffer comprising 12mM CaCl₂; 12mM MgCl₂; 200mM NaCl; 20mM Tris-HCl, pH 8.0; and 1mM EDTA.

The restriction enzymes EcoRI and NcoI were used in the high salt buffer and were obtained from New England Biolabs, Beverly, Massachusetts. The restriction enzymes BglII, BamHI, HindIII and PstI were used in medium salt buffer and were obtained from New England Biolabs, Beverly, Massachusetts. The restriction enzyme HpaI was used in the low salt buffer. The DNA ligase was used in ligase buffer and was obtained from New England Biolabs, Beverly, Massachusetts. The nuclease BAL31 was used in BAL31 nuclease buffer and was obtained from Bethesda Research Laboratories, Gaithersburg, Maryland. Bovine serum albumin was also obtained from Bethesda Research Laboratories.

Ampicillin, kanamycin sulfate, chloramphenicol and tetracycline were obtained from Sigma Chemical Company (Sigma), St. Louis, Missouri. Desthiobiotin was also obtained from Sigma Chemical Company. Biotin was obtained from either Sigma or from J. T. Baker Chemical Company, Phillipsburg, New Jersey. Strain BM4062 which has been deposited (August 23, 1985) with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland as A.T.C.C. 53238, had lesions in the *bioR* function and the *bioA* function. The *bioR* mutants were temperature sensitive in that they were viable at low temperatures (-28°C .) but unable to grow at high temperatures (-43°C). Depending on the specific mutant in question, the lethal effect of high temperature could be reversed by adding exogenous biotin. The other plasmids and strains employed in the Examples are summarized in Table II. Except for SA291, all of the *bioR*⁻ strains listed in Table II are reported in Barker et al., J. Bacteriol., 143, 789-800 (1980) or Campbell, et al., Proc. Natl. Acad. Sci. (USA), 69, 676-680 (1972) wherein they are all described as biotin requiring strains. Strain SA291 has been reported in Cleary, et al., J. Bacteriol., 112, 830-839 (1972). In Table II, it should also be noted that *BioR*⁺ refers to a genotype giving rise to a "partially defective" *BioR* gene product.

The following assays were utilized to determine the biotin concentration of the samples in the following Examples.

45 Microbiological Assay

The biotin concentration was determined by "cross-feeding" SA291 cells with the biotin produced by the designated strain. Initially, SA291 was cultured overnight in GMH broth (9 g/L vitamin assay Casamino Acids (Difco, Detroit, Michigan); 4 g/L glucose; 20 $\mu\text{g/L}$ 1-histidine; 40 $\mu\text{g/L}$ thiamine; 1mM MgSO₄; 6 g/L Na₂HPO₄; 3 g/L KH₂PO₄; 0.5 g/L NaCl; and 1 g/L NH₄Cl) supplemented with 300 pM d-biotin (50 ml volume) at 37°C. The overnight culture was diluted 400 fold in GMH broth. 2 mL samples of diluted culture were dispensed into test tubes. Varying concentrations of the sample to be analyzed were added to the tubes. The assay was standardized by adding one each of the following concentrations of d-biotin (Sigma Chemical Company, St. Louis, Missouri) to six individual tubes: 30 pM, 100 pM, 300 pM, 1000 pM, 3000 pM added d-biotin. All tubes were cultured overnight at 37°C. The optical density of resulting cultures were determined and the unknowns were correlated with the standards.

Spectrophotometric Assay

The biotin concentration was determined by a biotin spectrophotometric assay derived from McCormick et al., *Analytical Biochemistry*, 34, 226-236 (1970). Specifically, 100 microliters of sample were transferred to a test tube to which 900 microliters of water were added. Concentrated H₂SO₄ (5 microliters) was added to reduce the pH of the solution to less than 2. To the solution was added 1 mL of n-butanol and the resulting solution was vortexed for one minute and then centrifuged for one minute at 3200 rpm. The upper, butanol phase of the solution was then transferred to a 1.5 mL Eppendorf tube. The solvent was evaporated to dryness before the addition of 100 microliters of 2% (v/v) sulfuric acid and 100 microliters of 0.2% (w/v) 4-dimethylaminocinnamaldehyde. The resulting solution was vortexed and then allowed to stand undisturbed for 5-10 minutes before the addition of 800 µl of ethanol. The absorbance (A) of the solution at 532 nm was determined on a Gilford Response spectrophotometer. From the absorbance, the biotin concentration (C) in moles/liter was obtained for the 1 cm path length from the equation:

$$C = (5.7 \times 10^{-5})A$$

EXAMPLE 5

The various cultures utilized in Examples 6 and 7 were constructed by transforming an appropriate host strain with a plasmid derived from Examples 1 or 2 accordance with the procedures described by Hanahan, supra. The host strains and plasmids utilized are listed in Table II.

EXAMPLE 625 Flask Method

A flask containing 30 mL of GMH broth was inoculated with a volume of a culture listed in Table II. The resulting cultures were incubated with shaking at 37° C. At 0 hr, 6 hr, 21 hr, 30 hr and 45 hr, an aliquot was removed and filter sterilized. (A 2.5 mL aliquot was removed at 0 hr, while 1 mL aliquots were removed at the other time points). The sterilized samples were assayed in accordance with the microbiological assay previously described and the results (Run Nos. 1-7) are illustrated in Table II.

EXAMPLE 735 Fermentor Method

The appropriate host bearing plasmids were cultured overnight in GMH broth. A 10 mL aliquot of the culture was added to 1000 mL of GMH broth supplemented with 20 mL of 1% alanine, 20 mL of 1% methionine, 20 mL of 0.7% cysteine and appropriate antibiotic (final concentration of 50mg/L ampicillin was used when plasmid pKA5 was employed and 25 mg/L of kanamycin sulfate was used when plasmid pKH4 was employed). The fermentation was performed in a New Brunswick Bio-Flo fermentor with the following features: 1) constant agitation at 600 rpm; 2) sparging with air; 3) pH control between 6.8 and 7.2 by automatic addition of concentrated ammonia hydroxide; 4) temperature control and 5) a slow feed during the course of the fermentation. Dissolved oxygen and carbon dioxide evolution were not monitored. The feed consisted of: 12% glucose; 0.6% vitamin assay casamino acids, 35 µM magnesium sulfate, 0.2% alanine, 0.2% methionine, 0.12% cysteine, 7 µM sodium molybdate and half strength M9 minimal salts. The feed was initiated eight hours after inoculation at a continuous rate of 14 mL/hr. The temperature at the time of inoculation was 30° C. When the optical density of the culture reached approximately 10, measured at 600 nanometers, the temperature was incrementally raised to 40° C. The biotin concentrations reported in Table II were determined 24 hours after inoculation using either the microbiological assay (Run Nos. 8, 9 and 10) or the spectrophotometric assay (Run Nos. 10 and 12).

TABLE II
Biotin Production Determined by Bioassay

Run No.	Host Strain	Relevant Genotype	Plasmid	Relevant Genotype	Fermentation Method	Concentration of Biotin
1	SA291	<u>bio⁻ bio⁺ birA⁺</u>	pBP5	<u>bio⁺</u>	Flask	200 nM (50 µg/L)
2	S965	<u>bio⁺ bio⁺ birA^{TS}</u>	pBP5	<u>bio⁺</u>	Flask	600 nM - 1000 nM 150-250 µg/L
3	BM4110	<u>bio⁻ bio⁻ birA⁺</u>	pBP5	<u>bio⁺</u>	Flask	200 nM (50 µg/L)
4	BM4086	<u>bio⁻ bio⁻ birA⁺</u>	pBP5	<u>bio⁺</u>	Flask	100 nM (25 µg/L)
5	BM4084	<u>bio⁻ bio⁻ birA⁺</u>	pBP5	<u>bio⁺</u>	Flask	300 nM (75 µg/L)
6	BM4062	<u>bio⁻ bio⁻ birA^{TS}</u>	pKH4	<u>bio⁺</u>	Flask	2-6 µM (0.5-1.5 mg/L)
7	BM4062	<u>bio⁻ bio⁻ birA^{TS}</u>	pKA5	<u>bio⁺</u>	Flask	2-6 µM (0.5-1.5 mg/L)
8	S965	<u>bio⁺ bio⁺ birA^{TS}</u>	pBP5	<u>bio⁺</u>	Fermentor	6-10 µM (1.5-2.5 mg/L)
9	BM4062	<u>bio⁻ bio⁻ birA^{TS}</u>	pKA5	<u>bio⁺</u>	Fermentor	80 µM (20 mg/L)
10	BM4062	<u>bio⁻ bio⁻ birA^{TS}</u>	pKA5	<u>bio⁺</u>	Fermentor	125 µM (30 mg/L)
11	BM4062	<u>bio⁻ bio⁻ birA^{TS}</u>	pKH4	<u>bio⁺</u>	Fermentor	50 µM (12 mg/L)
12	BM4062	<u>bio⁻ bio⁻ birA^{TS}</u>	pKH4	<u>bio⁺</u>	Fermentor	80 µM (20 mg/L)

Although differences among assays for biotin make a direct comparison difficult, a comparison of the results reported in Table II with medium concentrations of biotin reported for wild type and mutant *E. coli* serve to illustrate the improvement in biotin production afforded by the present invention. For example, according to Campbell, et al., *Proc. Natl. Acad. Sci. (USA)*, 69, 676-680 (1972), the media concentration of biotin for wild type *E. coli* is less than 0.05nM and for strain S942 (a *birA⁻* strain) is 30-90nM. According to Pai, J. Bacteriol., 112, 1280-1287 (1972), for a strain P48, which may be employed as a biotin retention deficient-mutant-strain according to the present invention, the media concentration of biotin is 1000-times that of the wild type.

EXAMPLE 7

Cultures used in this Example are described in Table III.

TABLE III

Culture			
Strain	Genotype	Plasmid	Genotype
FM6	bio ⁺ , bioR ⁺ , birA ⁺	None	-
FM6	bio ⁺ , bioR ⁺ , birA ⁺	pKA5	bio ⁺
S965	bio ⁺ , bioR ^P , birA ^{TS}	None	-
S965	bio ⁺ , bioR ^P , birA ^{TS}	pKA5	bio ⁺
BM4062	bio ⁻ , bioR ⁻ , birA ^{TS}	None	-
BM4062	bio ⁻ , bioR ⁻ , birA ^{TS}	pKA5	bio ⁺

Each culture was inoculated into Luria broth (1.0% casamino acids, 0.5% yeast extracts, 0.5% sodium chloride) and incubated overnight at 30 °C. The optical density of each culture was determined and represented in Table IV. A ten-fold dilution was necessary due to density of the culture.

TABLE IV

Culture				
Strain	Plasmid	A ⁶⁰⁰ /10	A ⁶⁰⁰	Inoculum
FM6	None	0.6205	6.205	5 µL
FM6	pKA5	0.6056	6.056	5 µL
BM4062	None	0.2919	2.919	10 µL
BM4062	pKA5	0.5768	5.768	5 µL
S965	None	0.4920	4.920	6 µL
S965	pKA5	0.4379	4.379	7 µL

A flask containing 30 mL of GMH broth was inoculated with the volume of the six cultures listed in the Table IV above. The resulting cultures were incubated with shaking at 37 °C. At 0 hr, 6 hr, 21 hr, 30 hr and 45 hr an aliquot was removed and filter sterilized. (A 2.5 mL aliquot was removed at 0 hr, while 1 mL aliquots were removed at the other time points). The sterilized samples were assayed in accordance with the microbiological assay previously described and the results are illustrated in Table V.

TABLE V

Biotin Concentration Time Dependence of Biotin Accumulation					
Culture	0-hr	6 hr	21-hr	30 hr	45-hr
Strain/Plasmid					
FM6	<0.5nM	(1)	<1.6nM	<1.6 nM	<1.6 nM
FM6/pKA5	<0.5nM	<1nM	12nM	17nM	6.7nM
S965	<1.6nM	13 nM	133 nM	140nM	137nM
S965/pKA5	<1.6 nM	43 nM	150 nM	173nM	143nM
BM4062	<0.5nM	<1.6nM	<1.6nM	<1.6nM	<1.6nM
BM4062/pKA5	<1.6nM	(2)	3.1 µM	3.2 µM	1.5 µM
(1) Time point was contaminated; no value obtainable.					
(2) No data point in the linear range of the assay.					

While the present invention has been described in terms of preferred embodiments, it is understood that variations and improvements will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations which come within the scope of the invention as claimed.

5

Claims

1. A system for the production of biotin comprising: a microorganism having a birA⁻ genotype and plasmid DNA, within said cell, encoding at least one gene product of the biotin operon or a functional homolog thereof.
2. The system as recited in Claim 1 wherein said microorganism has a birA^{ts} genotype.
3. The system as recited in Claim 1 or 2 wherein said plasmid DNA encodes the biotin operon.
4. The system as recited in Claim 3 wherein said microorganism has a bioR⁻ genotype.
5. The system as recited in Claim 3 wherein said microorganism has a bio⁻ genotype.
6. The system as recited in any one of Claims 1 to 5 wherein said plasmid DNA is uvrB⁻.
7. The system as recited in any one of Claims 1 to 6 wherein said plasmid DNA encodes biotin synthetase or a functional homolog thereof.
8. A system for the production of biotin comprising: a microorganism having a (bio⁻, birA⁻, bioR⁻) genotype; and plasmid DNA, within said cell, wherein said plasmid DNA encodes the biotin operon and wherein said plasmid DNA is uvrB⁻.
9. A system for the production of biotin comprising: a microorganism having a (bio⁻, birA^{ts}, bioR⁻) genotype; and plasmid DNA, within said cells, wherein said plasmid DNA encodes the biotin operon and wherein said plasmid DNA is uvrB⁻.
10. A method for the enhanced production of biotin which comprises transforming a microorganism having a birA⁻ genotype with a plasmid DNA encoding a gene product of the biotin operon or a functional homolog thereof.
11. A method of converting desthiobiotin to biotin comprising the steps of: culturing a host microorganism, having a birA⁻ genotype and having plasmid DNA encoding a bioB gene product of the biotin operon or a functional homolog thereof, in a medium containing desthiobiotin.

40

Patentansprüche

1. System Für die Produktion von Biotin, welches umfaßt: einen Mikroorganismus mit einem birA⁻-Genotyp und Plasmid-DNA, in besagter Zelle, die wenigstens ein Genprodukt des Biotin-Operons oder ein funktionelles Homolog desselben codiert.
2. System nach Anspruch 1, dadurch gekennzeichnet, daß besagter Mikroorganismus einen birA^{ts}-Genotyp hat.
3. System nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß besagte Plasmid-DNA das Biotin-Operon codiert.
4. System nach Anspruch 3, dadurch gekennzeichnet, daß besagter Mikroorganismus einen bioR⁻-Genotyp hat.
5. System nach Anspruch 3, dadurch gekennzeichnet, daß besagter Mikroorganismus einen bio⁻-Genotyp hat.

55

6. System nach einem der Ansprüche 1 bis 5, dadurch gekennzeichnet, daß besagte Plasmid-DNA uvrB⁻ ist.
7. System nach einem der Ansprüche 1 bis 6, dadurch gekennzeichnet, daß besagte Plasmid-DNA Biotin-Synthetase oder ein funktionelles Homolog derselben codiert.
8. System für die Produktion von Biotin, welches umfaßt: einen Mikroorganismus mit einem (bio⁻, birA⁻, bioR⁻)-Genotyp; und Plasmid-DNA, in besagter Zelle, wobei besagte Plasmid-DNA das Biotin-Operon codiert und wobei besagte Plasmid-DNA uvrB⁻ ist.
9. System für die Produktion von Biotin, welches umfaßt: einen Mikroorganismus mit einem (bio⁻, birA^{ts}, bioR⁻)-Genotyp; und Plasmid-DNA, in besagten Zellen, wobei besagte Plasmid-DNA das Biotin-Operon codiert und wobei besagte Plasmid-DNA uvrB⁻ ist.
10. Verfahren zur gesteigerten Produktion von Biotin, welches umfaßt, daß ein Mikroorganismus mit einem birA⁻-Genotyp mit einer Plasmid-DNA, die ein Genprodukt des Biotin-Operons oder ein funktionelles Homolog desselben codiert, transformiert wird.
11. Verfahren zur Umwandlung von Desthiobiotin zu Biotin, welches die Schritte umfaßt: Kultivieren eines Wirtsmikroorganismus, mit einem birA⁻-Genotyp und mit Plasmid-DNA, die ein bioB-Genprodukt des Biotin-Operons oder eines funktionelles Homolog desselben codiert, in einem Medium, das Desthiobiotin enthält.

Revendications

1. Système de production de biotine comprenant: un microorganisme ayant un génotype birA⁻ et de l'ADN plasmidique, dans ladite cellule, qui code pour au moins un produit génétique de l'opéron biotine ou pour un de ses homologues fonctionnels.
2. Système selon la revendication 1, dans lequel ledit microorganisme a un génotype birA^{ts}.
3. Système selon la revendication 1 ou 2, dans lequel ledit ADN plasmidique code pour l'opéron biotine.
4. Système selon la revendication 3, dans lequel ledit microorganisme a un génotype bioR⁻.
5. Système selon la revendication 3, dans lequel ledit microorganisme a un génotype bio⁻.
6. Système selon l'une quelconque des revendications 1 à 5, dans lequel ledit ADN plasmidique est uvrB⁻.
7. Système selon l'une quelconque des revendications 1 à 6, dans lequel ledit ADN plasmidique code pour la biotine synthétase ou pour un de ses homologues fonctionnels.
8. Système pour la production de biotine, comprenant: un microorganisme ayant un génotype (bio⁻, birA⁻, bioR⁻); et un ADN plasmidique, dans ladite cellule, dans lequel ledit ADN plasmidique code pour l'opéron biotine et dans lequel ledit ADN plasmidique est uvrB⁻.
9. Système pour la production de biotine, comprenant: un microorganisme ayant un génotype (bio⁻, birA^{ts}, bioR⁻); et un ADN plasmidique, dans lesdites cellules, dans lequel ledit ADN plasmidique code pour l'opéron biotine et dans lequel ledit ADN plasmidique est uvrB⁻.
10. Méthode de production renforcée de biotine, qui comprend la transformation d'un microorganisme ayant un génotype birA⁻ avec un ADN plasmidique codant pour un produit génétique de l'opéron biotine ou pour un de ses homologues fonctionnels.
11. Méthode de conversion de desthiobiotine en biotine, qui comprend les étapes selon lesquelles on cultive un microorganisme hôte ayant un génotype birA⁻ et ayant de l'ADN plasmidique codant pour un produit du gène bioB de l'opéron biotine ou pour un de ses homologues fonctionnels, dans un

milieu contenant de la desthiobiotin .

5

10

15

20

25

30

35

40

45

50

55




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig.7




Fig. 8




Fig. 9